

Background characteristics of respondents**Percent distribution of background characteristics of women age 15 to 49**

Background characteristics	Urban	Total	Percentage of sample	Weighted number	Unweighted number
Age group					
15–19	100.0	100.0	14.6	299	212
20–24	100.0	100.0	12.4	256	180
25–29	100.0	100.0	18.0	371	271
30–34	100.0	100.0	18.4	378	266
35–39	100.0	100.0	17.1	351	230
40–44	100.0	100.0	11.9	244	165
45–49	100.0	100.0	7.7	159	108
Marital status					
Married	100.0	100.0	58.0	1,193	827
Living together	100.0	100.0	3.6	73	56
Divorced	100.0	100.0	3.2	67	42
Widowed	100.0	100.0	2.1	43	26
Never married	100.0	100.0	33.0	679	479
Parity					
None	100.0	100.0	35.4	729	507
1-2	100.0	100.0	31.3	644	444
3-4	100.0	100.0	25.4	523	368
5 or more	100.0	100.0	7.8	161	112
Education					
Never attended	100.0	100.0	2.5	52	36
Primary	100.0	100.0	11.2	230	161
Secondary	100.0	100.0	53.4	1,098	796
Higher	100.0	100.0	32.9	676	438
Wealth quintile					
Lowest	100.0	100.0	16.6	341	259
Lower	100.0	100.0	19.2	395	291
Middle	100.0	100.0	22.0	453	308
Higher	100.0	100.0	21.7	447	309
Highest	100.0	100.0	20.5	422	265
Total	100.0	100.0	100.0	2,057	1,432

Current use of contraception by background characteristics**Percentage of women ages 15 to 49 currently using a contraceptive method, by type and background characteristics**

Background characteristic	Any method		Modern method ¹		Traditional CPR	
	All women	Married women ²	All women	Married women ²	All women	Married women ²
Number	2,057	1,267	2,057	1,267	2,057	1,267
Total	26.4	32.5	19.7	22.9	6.7	9.6
Age group						
15–19	2.3	(0.0)	1.5	(0.0)	0.7	(0.0)
20–24	27.7	25.2	24.2	18.3	3.5	6.9
25–29	32.1	31.6	26.0	24.6	6.1	7.0
30–34	28.0	31.3	20.7	23.5	7.3	7.7
35–39	38.3	39.8	26.3	25.8	12.1	14.0
40–44	31.4	34.8	21.6	23.3	9.8	11.4
45–49	18.8	23.4	11.9	14.9	6.9	8.5
Marital status						
Married ²	32.5	na	22.9	na	9.6	na
Unmarried, sexually active	43.0	na	38.5	na	4.5	na
Parity						
0–1	16.4	16.2	13.0	10.6	3.4	5.6
2–3	33.2	34.4	24.9	25.5	8.3	8.9
4 or more	41.4	45.1	28.6	30.4	12.8	14.7
Education						
Never attended	8.3	11.8	5.7	8.0	2.7	3.8
Primary	20.3	25.1	17.4	21.4	2.9	3.8
Secondary	23.8	31.1	17.6	21.6	6.2	9.5
Higher	34.2	39.4	25.0	26.9	9.2	12.6
Wealth quintile						
Lowest	16.0	19.4	11.7	15.8	4.3	3.5
Lower	23.9	26.3	20.5	20.8	3.4	5.5
Middle	25.7	31.5	20.1	22.7	5.6	8.7
Higher	28.9	34.1	21.2	23.5	7.7	10.7
Highest	35.2	47.4	23.3	29.8	11.9	17.7

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

²Married includes married women and those who are living with a man

na = not applicable

() = Estimate generated from less than 25 unweighted observations

Contraceptive method mix by background characteristics

Percent distribution of current contraceptive users ages 15 to 49, by method type, marital status and background characteristics

All users										
N = 544										
Background characteristics	Sterilization	Implants	IUD	Injectables	Pill	Emergency contraception	Condom	Other modern	Other traditional	Total
Total	1.1	3.8	5.1	11.1	13.4	8.2	30.2	1.6	25.5	100.0
Age group										
15–19	(0.0)	(0.0)	(0.0)	(0.0)	(30.9)	(0.0)	(36.6)	(0.0)	(32.5)	(100.0)
20–24	0.0	0.0	0.0	1.0	7.1	16.7	62.4	0.0	12.7	100.0
25–29	0.0	2.2	2.4	10.0	20.2	7.1	39.0	0.0	19.0	100.0
30–34	2.2	4.3	11.7	17.7	6.9	6.8	20.3	3.9	26.2	100.0
35–39	0.6	5.4	3.7	15.1	14.4	9.5	17.4	2.4	31.4	100.0
40–44	2.6	6.2	2.4	11.1	12.9	4.5	29.0	0.0	31.2	100.0
45–49	(2.7)	(4.1)	(19.9)	(0.0)	(16.8)	(2.6)	(13.3)	(4.1)	(36.5)	(100.0)
Marital status										
Married ¹	1.3	4.5	6.4	14.0	13.9	4.7	23.7	2.1	29.5	100.0
Unmarried, sexually active	0.0	0.0	2.7	1.6	14.6	19.8	50.8	0.0	10.5	100.0
Parity										
0–1	0.5	0.6	0.6	2.0	9.4	13.0	53.2	0.0	20.8	100.0
2–3	1.6	3.3	5.8	12.0	17.8	6.6	24.7	3.1	25.1	100.0
4 or more	1.0	7.6	9.0	19.4	11.7	5.4	14.0	1.1	30.9	100.0
Education										
Never attended	(0.0)	(0.0)	(35.2)	(0.0)	(32.8)	(0.0)	(0.0)	(0.0)	(32.0)	(100.0)
Primary	0.0	3.8	4.4	20.6	27.7	5.2	21.0	2.8	14.5	100.0
Secondary	0.0	4.3	4.2	12.2	12.3	8.5	31.3	1.1	26.1	100.0
Higher	2.6	3.2	5.8	8.1	11.4	8.6	31.5	1.8	26.9	100.0
Wealth quintile										
Lowest	0.0	1.5	0.0	17.0	27.5	6.7	20.5	0.0	26.9	10.0
Lower	0.0	4.2	2.7	8.4	10.5	11.4	48.6	0.0	14.3	100.0
Middle	0.0	4.6	4.9	9.2	20.2	6.5	31.5	1.1	21.9	100.0
Higher	0.0	2.9	6.7	16.5	12.0	6.6	27.6	0.9	26.7	100.0
Highest	4.0	4.4	7.4	7.4	6.0	9.5	23.4	4.1	33.9	100.0

¹Married includes married women and those who are living with a man

na = not applicable

() = Estimate generated from less than 25 unweighted observations


() = Estimate generated from less than 5 unweighted observations

Married users										
N = 412										
Background characteristics	Sterilization	Implants	IUD	Injectables	Pill	Emergency contraception	Condom	Other modern	Other traditional	Total
Total	1.3	4.5	6.4	14.0	13.9	4.7	23.7	2.1	29.5	100.0
Age group										
15–19	na	na	na	na	na	na	na	na	na	na
20–24	0.0	0.0	0.0	3.9	3.9	10.9	53.8	0.0	27.5	(100.0)
25–29	0.0	3.1	2.4	14.2	21.5	5.3	31.3	0.0	22.2	100.0
30–34	1.6	4.8	12.0	18.7	7.7	5.7	20.4	4.3	24.7	100.0
35–39	0.7	4.7	4.3	17.3	15.0	4.9	15.1	2.8	35.1	100.0
40–44	3.0	7.1	2.7	10.2	12.9	1.2	30.0	0.0	32.9	100.0
45–49	(2.7)	(4.1)	(19.9)	(0.0)	(16.8)	(2.7)	(13.3)	(4.1)	(36.5)	(100.0)
Marital status										
Married ¹	na	na	na	na	na	na	na	na	na	na
Unmarried, sexually active	na	na	na	na	na	na	na	na	na	na
Parity										
0–1	0.0	1.9	0.0	6.1	5.9	3.8	48.0	0.0	34.4	100.0
2–3	1.7	2.6	5.7	11.3	17.3	6.2	26.0	3.3	25.9	100.0
4 or more	1.1	8.0	9.5	20.4	12.3	2.9	12.1	1.2	32.5	100.0
Education										
Never attended	(0.0)	(0.0)	(35.2)	(0.0)	(32.8)	(0.0)	(0.0)	(0.0)	(32.0)	(100.0)
Primary	0.0	3.9	4.5	21.4	28.7	5.4	18.1	2.9	15.0	100.0
Secondary	0.0	4.9	4.7	15.0	15.1	6.2	21.8	1.5	30.7	100.0
Higher	3.1	4.3	8.0	11.2	8.2	2.8	28.0	2.5	31.9	100.0
Wealth quintile										
Lowest	0.0	2.2	0.0	25.5	37.0	3.6	13.3	0.0	18.3	100.0
Lower	0.0	6.1	3.9	12.3	15.3	7.2	34.2	0.0	20.9	100.0
Middle	0.0	4.0	5.5	10.9	20.5	6.7	23.1	1.5	27.7	100.0
Higher	0.0	3.7	7.6	19.4	8.6	4.5	23.9	1.2	31.2	100.0
Highest	4.3	5.4	9.2	9.1	6.0	2.4	21.5	5.0	37.2	100.0

() = Estimate generated from less than 25 unweighted observations

() = Estimate generated from less than 5 unweighted observations

Current Method Mix Among All and Married Contraceptive Users (Age 15-49)



Unmet need for family planning

Percentage of women age 15 to 49 with unmet need for family planning, by marital status and background characteristics

Background characteristics	All women			Married women ¹		
	2,057			1,267		
Number	For spacing	For limiting	Total	For spacing	For limiting	Total
Total	10.0	5.6	15.6	12.7	8.8	21.5
Age group						
15–19	6.1	0.3	6.4	(72.6)	(0.0)	(72.6)
20–24	9.3	2.0	11.3	10.3	6.8	17.1
25–29	13.8	0.8	14.6	17.2	1.2	18.3
30–34	14.1	7.8	21.9	16.2	9.2	25.4
35–39	11.8	6.1	18.0	13.0	7.3	20.3
40–44	6.5	12.8	19.3	6.3	15.4	21.7
45–49	1.5	14.8	16.3	1.9	18.4	20.3
Marital status						
Married ¹	12.7	8.8	21.5	na	na	na
Unmarried, sexually active	27.5	2.3	29.8	na	na	na
Parity						
0–1	6.6	0.4	7.0	9.2	1.0	10.2
2–3	16.2	7.8	24.0	16.5	8.3	24.9
4 or more	8.9	15.2	24.2	9.4	17.1	26.5
Education						
Never attended	5.5	7.9	13.4	5.2	11.2	16.3
Primary	11.3	8.4	19.8	14.0	10.8	24.8
Secondary	11.8	5.8	17.5	15.3	9.6	24.8
Higher	7.2	4.1	11.3	9.1	6.5	15.6
Wealth quintile						
Lowest	17.2	6.4	23.6	23.1	10.2	33.2
Lower	11.2	6.0	17.2	14.6	9.7	24.3
Middle	11.2	4.0	15.1	14.9	6.4	21.3
Higher	7.6	7.2	14.8	7.9	10.5	18.4
Highest	4.6	4.5	9.1	6.4	7.4	13.9

¹Married includes married women and those who are living with a man

na = not applicable

() = Estimate generated from less than 25 unweighted observations

Need and demand for family planning

Percentage of women age 15 to 49 with met need, unmet need, and total demand for family planning, and percentage of demand for family planning satisfied with modern methods, by marital status and background characteristics

Background characteristics Number	All women 2,057					Married women ² 1,267				
	Any Method	Modern method ¹	Unmet need	Total demand	Demand satisfied by modern method	Any Method	Modern method ¹	Unmet need	Total demand	Demand satisfied by modern method
Total	26.4	19.7	15.6	42.0	46.8	32.5	22.9	21.5	54.0	42.5
Age group										
15–19	2.3	1.5	6.4	8.7	17.6	(0.0)	(0.0)	(72.6)	(72.6)	(0.0)
20–24	27.7	24.2	11.3	39.0	62.1	25.2	18.3	17.1	42.3	43.2
25–29	32.1	26.0	14.6	46.7	55.6	31.6	24.6	18.3	49.9	49.2
30–34	28.0	20.7	21.9	49.9	41.5	31.3	23.5	25.4	56.7	41.5
35–39	38.3	26.3	18.0	56.3	46.7	39.8	25.8	20.3	60.0	43.0
40–44	31.4	21.6	19.3	50.7	42.6	34.8	23.3	21.7	56.5	41.3
45–49	18.8	11.9	16.3	35.1	34.0	23.4	14.9	20.3	43.7	34.0
Marital status										
Married ²	32.5	22.9	21.5	54.0	42.5	na	na	na	na	na
Unmarried, sexually active	43.0	38.5	29.8	72.8	52.8	na	na	na	na	na
Parity										
0–1	16.4	13.0	7.0	23.3	55.6	16.2	10.6	10.2	26.4	40.2
2–3	33.2	24.9	24.0	57.2	43.5	34.4	25.5	24.9	59.3	43.0
4 or more	41.4	28.6	24.2	65.6	43.7	45.1	30.4	26.5	71.6	42.5
Education										
Never attended	8.3	5.7	13.4	21.8	26.1	11.8	8.0	16.3	28.1	28.5
Primary	20.3	17.4	19.8	40.1	43.3	25.1	21.4	24.8	49.9	42.8
Secondary	23.8	17.6	17.5	41.3	42.6	31.1	21.6	24.8	55.9	38.5
Higher	34.2	25.0	11.3	45.5	54.9	39.4	26.9	15.6	55.1	48.8
Wealth quintile										
Lowest	16.0	11.7	23.6	39.6	29.6	19.4	15.8	33.2	52.6	30.1
Lower	23.9	20.5	17.2	41.1	49.9	26.3	20.8	24.3	50.5	41.1
Middle	25.7	20.1	15.1	40.9	49.2	31.5	22.7	21.3	52.8	43.1
Higher	28.9	21.2	14.8	43.7	48.5	34.1	23.5	18.4	52.5	44.7
Highest	35.2	23.3	9.1	44.3	52.5	47.4	29.8	13.9	61.3	48.6

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

²Married includes married women and those who are living with a man

na = not applicable

() = Estimate generated from less than 25 unweighted observations

Pregnancy intentions

Unintended births among women age 15 to 49 with one or more births in the previous 5 years or who are currently pregnant: Percent distribution reporting last or current pregnancy was wanted then, wanted later or not wanted at all, by marital status and background characteristics

Background characteristics	All women with 1 or more births				Married women with 1 or more births			
	857				804			
Number	Wanted then	Wanted later	Wanted no more	Total	Wanted then	Wanted later	Wanted no more	Total
Total	76.7	15.9	7.4	100.0	78.1	14.8	7.1	100.0
Age group								
15–19	(24.8)	(44.6)	(30.6)	(100.0)	(33.0)	(48.3)	(18.7)	(100.0)
20–24	63.3	20.6	16.1	100.0	75.0	12.2	12.8	100.0
25–29	81.2	18.0	0.8	100.0	82.0	17.1	0.9	100.0
30–34	79.4	17.0	3.5	100.0	78.8	17.5	3.6	100.0
35–39	81.7	8.3	10.0	100.0	81.7	7.5	10.8	100.0
40–44	67.4	14.4	18.2	100.0	67.8	13.2	19.1	100.0
45–49	(42.5)	(36.9)	(20.6)	(100.0)	(42.5)	(36.9)	(20.6)	(100.0)
Marital status								
Married ¹	78.1	14.8	7.1	100.0	na	na	na	na
Unmarried, sexually active	(34.1)	(57.5)	(8.4)	(100.0)	na	na	na	na
Parity								
0–1	75.3	18.3	6.5	100.0	80.7	15.3	4.0	100.0
2–3	85.7	13.7	0.6	100.0	85.9	13.5	0.7	100.0
4 or more	59.5	17.9	22.6	100.0	58.3	17.1	24.6	100.0
Education								
Never attended	(100.0)	(0.0)	(0.0)	(100.0)	(100.0)	(0.0)	(0.0)	(100.0)
Primary	75.8	11.2	13.0	100.0	76.0	11.4	12.6	100.0
Secondary	72.1	17.8	10.0	100.0	74.6	15.5	9.9	100.0
Higher	82.1	15.9	2.0	100.0	82.3	15.8	1.8	100.0
Wealth quintile								
Lowest	74.3	20.5	5.3	100.0	76.2	18.5	5.4	100.0
Lower	75.8	14.6	9.5	100.0	78.9	12.9	8.2	100.0
Middle	76.7	17.2	6.1	100.0	76.9	16.7	6.3	100.0
Higher	81.3	9.4	9.3	100.0	82.2	8.3	9.5	100.0
Highest	74.5	18.8	6.7	100.0	75.8	18.4	5.8	100.0

¹Married includes married women and those who are living with a man

na = not applicable

() = Estimate generated from less than 25 unweighted observations

Contraceptive choice

Percent distribution of women age 15 to 49 who are currently using modern contraception¹ reporting on who decided on the contraceptive method, by background characteristics

Background characteristics	Self	Jointly with partner or provider	Provider alone/ partner alone/other	Total
Number	157	317	62	536
Total	29.3	59.1	11.6	100.0
Age group				
15–19	(32.5)	(30.9)	(36.6)	(100.0)
20–24	18.3	56.9	24.8	100.0
25–29	29.8	58.4	11.8	100.0
30–34	34.1	52.6	13.3	100.0
35–39	25.5	69.6	4.9	100.0
40–44	37.4	55.0	7.5	100.0
45–49	(30.8)	(60.4)	(8.8)	(100.0)
Marital status				
Married ²	27.9	61.4	10.7	100.0
Unmarried, sexually active	31.2	48.8	19.9	100.0
Parity				
0–1	27.4	55.3	17.3	100.0
2–3	29.3	60.8	9.9	100.0
4 or more	31.2	60.8	8.0	100.0
Education				
Never attended	68.0	0.0	32.0	100.0
Primary	30.8	56.6	12.7	100.0
Secondary	34.5	53.7	11.8	100.0
Higher	22.1	67.2	10.7	100.0
Wealth quintile				
Lowest	(46.5)	(40.3)	(13.2)	(100.0)
Lower	33.7	48.2	18.1	100.0
Middle	27.5	60.7	11.8	100.0
Higher	26.5	67.4	6.1	100.0
Highest	23.8	65.0	11.2	100.0

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

²Married includes married women and those who are living with a man

na = not applicable

() = Estimate generated from less than 25 unweighted observations

() = Estimate generated from less than 5 unweighted observations

Payment for family planning

Percentage of women age 15 to 49 currently using a modern contraceptive method¹ who paid fees for services, by marital status and background characteristics

Background characteristics	All modern contraceptive users ¹	Married ² modern contraceptive users ¹
Number	544	412
Total	37.0	38.0
Age group		
15–19	(56.7)	(0.0)
20–24	35.4	56.4
25–29	44.5	49.5
30–34	32.8	33.3
35–39	37.6	38.2
40–44	30.8	26.8
45–49	(34.5)	(34.5)
Marital status		
Married ²	38.0	na
Unmarried, sexually active	37.8	na
Parity		
0–1	33.8	37.8
2–3	37.3	37.4
4 or more	39.8	39.0
Education		
Never attended	(0.0)	(0.0)
Primary	54.2	56.2
Secondary	35.0	36.5
Higher	36.5	35.9
Wealth quintile		
Lowest	54.8	71.3
Lower	34.9	32.4
Middle	42.4	42.7
Higher	31.9	33.8
Highest	32.0	31.2

() = Estimate generated from less than 25 unweighted observations

(/) = Estimate generated from less than 5 unweighted observations

Method Information Index Components

Percentage of current modern¹ contraceptive users who reported whether provider informed them about other methods, side effects and, if informed of side effects, what to do, by marital status and background characteristics²

Background characteristics	Informed about other methods		Informed about side effects		Told what to do if experienced side effects	
	All users	Married users ³	All users	Married users ³	All users	Married users ³
Number	544	412	404	289	135	112
Total	42.2	50.2	33.5	38.7	75.4	77.7
Age group						
15–19	(30.9)	(0.0)	(0.0)	na	na	na
20–24	14.5	32.6	15.7	(22.4)	(22.6)	(20.9)
25–29	38.0	47.2	23.1	25.9	(81.9)	(76.3)
30–34	55.6	58.9	46.1	47.6	88.4	(87.7)
35–39	42.7	47.6	39.7	44.0	(66.3)	(73.0)
40–44	51.8	49.6	40.2	32.9	(79.7)	(71.0)
45–49	(52.9)	(52.9)	(55.9)	(55.9)	(91.2)	(91.2)
Marital status						
Married ³	50.2	na	38.7	na	77.7	na
Unmarried, sexually active	15.6	na	29.2	na	62.5	na
Parity						
0–1	23.4	48.6	14.8	18.8	(60.8)	(60.6)
2–3	47.4	47.4	34.4	32.6	78.9	83.2
4 or more	54.6	54.5	54.5	54.3	77.1	75.2
Education						
Never attended	(35.2)	(35.2)	(51.8)	(51.8)	(100.0)	(100.0)
Primary	33.7	34.9	27.1	28.3	(57.6)	(57.6)
Secondary	46.5	52.6	36.9	40.3	75.2	73.9
Higher	39.2	51.8	30.8	39.9	78.6	86.1
Wealth quintile						
Lowest	25.9	33.5	22.7	(25.4)	(47.5)	(57.1)
Lower	33.3	41.2	32.5	31.1	(76.6)	(67.4)
Middle	42.2	47.9	24.4	30.3	(74.7)	(80.3)
Higher	44.0	53.4	46.8	53.8	82.7	(90.2)
Highest	52.4	58.9	34.3	41.9	(72.9)	(71.1)

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

²"Informed about other methods" asked of all current contraceptive users, "informed about side effects" asked of current modern contraceptive users, and "told what to do if experienced side effects" asked of current modern contraceptive users who indicated they were told about side effects.

³Married includes married women and those who are living with a man

na = not applicable

() = Estimate generated from less than 25 unweighted observations

() = Estimate generated from less than 5 unweighted observations

Perceived quality of care for family planning

Percentage of current modern¹ contraceptive users who would return, would refer a friend/relative, or would return and refer friend/relative to provider, by background characteristics

Background characteristics	Percent who would return to provider	Percent who would refer relative/friend	Percent who would return and refer friend/relative
Number	402	402	402
Total	77.8	75.5	70.2
Age group			
15–19	(100.0)	(93.8)	(93.8)
20–24	53.8	53.1	48.0
25–29	81.6	81.2	76.3
30–34	83.5	75.9	73.3
35–39	78.4	76.1	68.8
40–44	82.8	82.6	73.2
45–49	(91.2)	(91.2)	(91.2)
Marital status			
Married ²	81.2	80.6	74.0
Unmarried, sexually active	69.0	61.3	61.3
Parity			
0–1	66.5	64.3	60.4
2–3	83.4	82.5	77.0
4 or more	83.2	78.8	72.3
Education			
Never attended	(51.8)	(51.8)	(51.8)
Primary	83.5	69.2	69.2
Secondary	80.4	75.6	71.0
Higher	73.9	77.3	69.9
Wealth quintile			
Lowest	88.3	64.5	62.5
Lower	82.3	81.3	79.6
Middle	79.1	72.8	71.8
Higher	74.3	83.0	70.1
Highest	72.0	70.3	64.2

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

²Married includes married women and those who are living with a man

na = not applicable

() = Estimate generated from less than 25 unweighted observations

() = Estimate generated from less than 5 unweighted observations

Knowledge of family planning

Percentage of women age 15 to 49 who reported receiving family planning information from a health worker in the past 12 months, by marital status and background characteristics

Background characteristics	Visited by health worker who discussed family planning		Visited facility and provider discussed family planning		Either visited by health worker or visited facility and discussed family planning	
	All women	Married women ¹	All women	Married women ¹	All women	Married women ¹
Number	2,057	1,267	2,057	1,267	2,057	1,267
Total	11.1	13.6	24.7	37.0	29.9	41.7
Age group						
15–19	8.9	0.0	2.0	26.8	10.4	26.8
20–24	4.3	6.4	17.9	48.2	19.8	48.2
25–29	13.5	15.5	38.0	50.9	41.6	53.2
30–34	16.8	18.4	39.3	46.3	45.7	52.7
35–39	11.4	12.5	31.5	36.4	35.2	39.7
40–44	7.9	9.3	18.8	19.7	23.9	26.1
45–49	11.2	12.6	6.3	7.8	14.9	17.3
Marital status						
Married ¹	13.6	na	37.0	na	41.7	na
Unmarried, sexually active	9.0	na	6.7	na	12.9	na
Parity						
0–1	8.4	11.0	13.5	34.6	18.8	38.3
2–3	14.1	15.2	39.4	42.3	44.5	47.9
4 or more	13.0	13.4	29.3	30.1	34.3	34.3
Education						
Never attended	10.5	14.8	5.9	8.3	11.6	16.3
Primary	10.3	10.7	19.6	23.1	25.2	28.5
Secondary	11.9	14.7	24.0	38.8	29.9	43.7
Higher	10.2	13.2	29.0	42.7	32.8	46.6
Wealth quintile						
Lowest	9.7	14.7	18.2	29.1	22.2	33.7
Lower	13.9	16.4	24.5	38.1	30.3	41.4
Middle	12.5	14.6	26.1	39.1	32.3	43.5
Higher	11.8	14.3	26.1	35.8	32.5	43.6
Highest	7.3	8.2	27.0	41.0	30.3	43.8

List of indicators for sampling errors, PMA2016/Lagos-Round 3

Variable	Estimation	Base population
Currently using a modern method	Proportion	All women ages 15–49 Married women ages 15–49
Currently using a traditional method	Proportion	All women ages 15–49 Married women ages 15–49
Currently using any contraceptive method	Proportion	All women ages 15–49 Married women ages 15–49
Currently using injectables	Proportion	All women ages 15–49 Married women ages 15–49
Currently using male condoms	Proportion	All women ages 15–49 Married women ages 15–49
Currently using implants	Proportion	All women ages 15–49 Married women ages 15–49
Chose method by self or jointly in past 12 months	Proportion	All women ages 15–49 Married women ages 15–49
Paid fees for family planning services in past 12 months	Proportion	All women ages 15–49 Married women ages 15–49
Informed by provider about other methods	Proportion	All women ages 15–49 Married women ages 15–49
Informed by provider about side effects	Proportion	All women ages 15–49 Married women ages 15–49
Satisfied with provider: Would return and refer friend/relative to provider	Proportion	All women ages 15–49 Married women ages 15–49
Visited by health worker who talked about family planning information in past 12 months	Proportion	All women ages 15–49 Married women ages 15–49

Characteristics of service delivery points**Number and percentage of service delivery points (SDPs) by background characteristics**

Background characteristics	Number	Percentage
Total	124	100.0
SDP type		
Hospital	25	20.2
Health center ¹	57	46.0
Pharmacy ²	40	32.3
Other	2	1.6
Sector		
Public	68	54.8
Private	56	45.2
Offers family planning		
No	8	6.5
Yes	116	93.5

¹Includes Health Post, Health Clinic, and Maternity Center

²Includes Chemist and Patent Medicine Vendors

Charging fees for family planning

Percentage of service delivery points (SDPs) that charge fees for family planning services, by sector and background characteristics, among SDPs offering family planning

Background characteristic	Sector	
	Public	Private
Number of SDPs	67	49
Total	7.5	67.3
SDP type		
Hospital	(20.0)	(100.0)
Health center ¹	3.8	(50.0)
Pharmacy ²	na	61.1

¹Includes Health Post, Health Clinic, and Maternity Center

²Includes Chemist and Patent Medicine Vendors

() = Estimate generated from less than 25 unweighted observations

() = *Estimate generated from less than 5 unweighted observations*

Contraceptive choice: Availability of at least 3 or 5 modern contraceptive methods

Percentage of public and private service delivery points (SDPs) offering at least 3 or 5 modern contraceptive methods¹, by sector and background characteristics, among all SDPs

Background characteristic	3 or more methods		5 or more methods	
	Public	Private	Public	Private
Number of SDPs	68	56	68	56
Total	95.6	60.7	88.2	16.1
SDP type				
Hospital	(100.0)	(80.0)	(93.3)	(40.0)
Health Center ²	94.3	(50.0)	86.8	(25.0)
Pharmacy ³	na	55.0	na	10.0
Other	na	(100.0)	na	(0.0)

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

²Includes Health Post, Health Clinic, and Maternity Center

³Includes Chemist and Patent Medicine Vendors

na = not applicable

() = Estimate generated from less than 25 unweighted observations

() = Estimate generated from less than 5 unweighted observations

Contraceptive Choice: Availability of modern contraception, by method

Percentage of service delivery points (SDPs) offering different types of contraceptive methods¹, by sector and method, among all SDPs

Public SDPs							
Background characteristic	Contraceptive method						
	Pill	Injectables	IUD	Implants	Male condom	Female sterilization	Emergency contraception
Number of SDPs = 68							
Total	97.1	95.6	88.2	86.8	97.1	7.4	20.6
SDP type							
Hospital	(100.0)	(100.0)	(100.0)	(100.0)	(100.0)	(33.3)	(33.3)
Health center ²	96.2	94.3	84.9	83.0	96.2	0.0	17.0
Pharmacy ³	na	na	na	na	na	na	na
Other	na	na	na	na	na	na	na

Private SDPs							
Background characteristic	Contraceptive method						
	Pill	Injectables	IUD	Implants	Male condom	Female sterilization	Emergency contraception
Number of SDPs = 56							
Total	73.2	37.5	17.9	14.3	80.4	7.1	48.2
SDP type							
Hospital	(80.0)	(90.0)	(70.0)	(50.0)	(60.0)	(40.0)	(50.0)
Health center ²	(50.0)	(50.0)	(50.0)	(50.0)	(50.0)	(0.0)	(25.0)
Pharmacy ³	72.5	20.0	0.0	0.0	90.0	0.0	50.0
Other	(100.0)	(100.0)	(50.0)	(50.0)	(50.0)	(0.0)	(50.0)

¹Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

²Includes Health Post, Health Clinic, and Maternity Center

³Includes Chemist and Patent Medicine Vendors

() = Estimate generated from less than 25 unweighted observations

() = Estimate generated from less than 5 unweighted observations

Contraceptive stock-outs, by method

Percentage of service delivery points (SDPs) stocked out of specific modern contraceptives¹ at time of survey or stocked out in the last 3 months, by sector and background characteristics, among SDPs that offer each method

Public SDPs						
Background characteristic	Contraceptive method					
	Pill	Injectables	IUD	Implants	Male condom	Emergency contraception
Number of SDPs	66	65	60	59	66	14
Total	13.6	12.3	3.3	27.1	10.6	(28.6)
SDP type						
Hospital	(6.7)	(13.3)	(6.7)	(46.7)	(0.0)	(40.0)
Health center ¹	15.7	12.0	2.2	20.5	13.7	(22.2)
Pharmacy ²	na	na	na	na	na	na
Other	na	na	na	na	na	na

Private SDPs						
Background characteristic	Contraceptive method					
	Pill	Injectables	IUD	Implants	Male condom	Emergency contraception
Number of SDPs	41	21	10	8	45	27
Total	19.5	(14.3)	(10.0)	(12.5)	15.6	25.9
SDP type						
Hospital	(37.5)	(22.2)	(14.3)	(20.0)	(16.7)	(60.0)
Health center ¹	(50.0)	(0.0)	(0.0)	(0.0)	(50.0)	(0.0)
Pharmacy ²	13.8	(0.0)	na	na	13.9	20.0
Other	(0.0)	(50.0)	(0.0)	(0.0)	(0.0)	(0.0)

¹Includes Health Post, Health Clinic, and Maternity Center

²Includes Chemist and Patent Medicine Vendors

() = Estimate generated from less than 25 unweighted observations

() = Estimate generated from less than 5 unweighted observations

Number of family planning visits, by method

Total number of visits and new clients contraceptive services, by method type and background characteristics, among advanced¹ service delivery points (SDPs) that offer each method

Contraceptive Methods														
	Sterilization		Injectable		IUD		Implants		Male condom		Pills		Emergency Contraception	
	Total women	Total men	Total	New	Total	New	Total	New	Total	New	Total	New	Total	New
Number of SDPs	9	8	78		70		67		75		78		21	
Total	3	1	2,019	415	579	174	569	371	3,695	1,305	661	277	139	126
Background characteristic														
SDP type														
Hospital	3	1	596	148	420	84	248	146	438	289	296	124	115	115
Health center ¹	na	na	1,418	264	159	90	321	225	3,257	1,016	359	151	24	11
Other	na	na	5	3	0	0	0	0	0	0	6	2	0	0
Sector														
Public	0	0	1,985	394	486	141	565	369	3,594	1,206	626	262	28	15
Private	3	1	34	21	93	33	4	2	101	99	35	15	111	111

¹Includes Health Post, Health Clinic, and Maternity Center